U.S. DEPARTMENT OF COMMERCE National Technical Information Service

AD-A024 258

ELECTRONIC ANGULAR MOMENTUM TRANSFER IN THE I (DOUBLETP(1/2) + NF (A SINGLET DELTA) SYSTEM

AEROSPACE CORPORATION

PREPARED FOR
SPACE AND MISSILE SYSTEMS ORGANIZATION

17 March 1976

REPORT SAMSO-TR-76-58

Electronic Angular Momentum Transfer in the $I(^2P_{1/2}) + NF(a^1\Delta)$ System

ADA 024258

JOHN M. HERBELIN and M. A. KWOK
Aerophysics Laboratory
Laboratory Operations
The Aerospace Corporation
El Segundo, Calif. 90245

17 March 1976

Interim Report


APPROVED FOR PUBLIC RELEASE:
DISTRIBUTION UNLIMITED

Prepared for

SPACE AND MISSILE SYSTEMS ORGANIZATION
AIR FORCE SYSTEMS COMMAND

Los Angeles Air Force Station
P.O. Box 92960, Worldway Postal Center
Los Angeles, Calif. 90009

REPRODUCED BY
NATIONAL TECHNICAL
INFORMATION SERVICE
U. S. DEPARTMENT OF COMMERCE
SPRINGFIELD VA. 22161



REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
REPORT NUMBER	GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
SAMSO-TR-76-58		
TITLE (and Subtitie)		5. TYPE OF REPURT & PERIOD COVERED
ELECTRONIC ANGULAR MOMENTUM		Interim
TRANSFER IN THE $I(^2P_{1/2}) + NI$	$F(a^{l} \Delta)$	
SYSTEM		6. PERFORMING ORG. REPORT NUMBER
AUTHOR(a)		TR -0076(6240-40)-4 8. CONTRACT OR GRANT NUMBER(*)
John M. Herbelin and Munson A.	Kwok	F04701-75-C-0076
PERFORMING ORGANIZATION NAME AND ADDRESS		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
The Aerospace Corporation		
El Segundo, Calif. 90245		
1. CONTROLLING OFFICE NAME AND AODRESS		12. REPORT DATE
Space and Missile Systems Organization		17 March 1976
Air Force Systems Command	ii Zatioii	13. NUMBER OF PAGES
Los Angeles, Calif. 90009		7
4. MONITORING AGENCY NAME & ADDRESS(If different	from Controlling Office)	15. SECURITY CLASS. (of this report)
		Unclassified
		15. DECLASSIFICATION/DOWNGRADING
6. DISTRIBUTION STATEMENT (of this Report)		
7. DISTRIBUTION STATEMENT (of the abetract entered in	n Block 20, If different fro	m Report)
8. SUPPLEMENTARY NOTES		
9. KEY WOROS (Continue on reverse side if necessary and	I identify by block number)
Nitrogen Fluoride		
Energy Transfer, Electronic Sta	tes of Nitrogen	Fluoride
Electronic Energy Transfer		
Iodine Atoms		
Atom-Molecule Energy Transfer O. ABSTRACT (Continue on reverse elde if necessary and	S Identify by black number	,
A rate coefficient at 298 K of >2 deduced for the electronic energ		mol ⁻¹ -sec ⁻¹ has been
$I(^{2}P_{1/2}) + NF$ ($a^{1}\Delta) \longrightarrow I(^{2}P_{3/2})$	$+ NF(b^{1}\Sigma^{+})$

This report was submitted by The Aerospace Corporation, El Segundo, CA 90245, under Contract F04701-75-C-0076 with the Space and Missile Systems Organization, Deputy for Advanced Space Programs, P.O. Box 92960, Worldway Postal Center, Los Angeles, CA 90009. It was reviewed and approved for The Aerospace Corporation by W. R. Warren, Jr., Director, Aerophysics Laboratory. Lt. Ronald C. Lawson, SAMSO/DYN, was the project officer.

This report has been reviewed by the Information Office (OI) and is releasable to the National Technical Information Service (NTIS). At NTIS, it will be available to the general public, including foreign nations.

This technical report has been reviewed and is approved for publication. Publication of this report does not constitute Air Force approval of the report's findings or conclusions. It is published only for the exchange and stimulation of ideas.

FOR THE COMMANDER

Ronald C. Lawson

1st Lt., United States Air Force Research Applications Directorate Deputy for Advanced Space Programs

ii

PREFACE

The authors acknowledge the assistance of R. H. Ueunten in the experiment.

CONTENTS

PREFACE		1
ELECTRO	NIC ANGULAR MOMENTUM TRANSFER IN THE	
I(² P _{1/2}) +	$NF(a^{1}\Delta)$ SYSTEM	
Figure 1.	Axial Variations of the Number Density of NF($b^1\Sigma^+$), Averaged Along the Tube Diameter	-

Preceding page blank

ELECTRONIC ANGULAR MOMENTUM TRANSFER IN THE $I(^2P_{1/2}) + NF(a^1\Delta)$ SYSTEM

Evidence has been observed for a nearly gaskinetic rate coefficient for the E-E transfer:

$$I(^{2}P_{1/2}) + NF(a^{1}\Delta) \rightarrow I(^{2}V_{3/2}) + NF(b^{1}\Sigma^{+}) + 110 \text{ cm}^{-1}$$
 (1)

by using the large-diameter, medium-pressure flow tube facility. Such a large rate coefficient had been anticipated by previous studies *, ***, 2 that describe the role of electronic spin conservation in the reactive F + H₂/H + NF₂ systems in producing NF electronically excited species. The experiment involves determination of the axial variations of NF(a $^1\Delta$) and NF(b $^1\Sigma^+$) number densities in a sequence of carefully selected reagent flow conditions. This is followed by observation of the (a $^1\Delta$ -X $^3\Sigma^-$) and (b $^1\Sigma^+$ -X $^3\Sigma^-$) spontaneous emissions in the axially translating reaction zone. The enhancement of NF(b $^1\Sigma^+$) by the presence of I($^2P_{1/2}$) can then be observed, and an estimate can be made of the rate coefficient by using a chemical kinetic model that fits the experimental number densities.

From the passage of Ar + NF₃ through an rf discharge, F and 1% NF₂ are produced, which can be reacted with D₂ and HI injected rapidly into the principal flow from an axially movable tee injector located on the tube centerline. The reaction F + D₂ yields D atoms in excess, which, in turn, react with NF₂ to predominantely produce. NF(a $^{1}\Delta$). The excess D also rapidly

Preceding page blank

J. M. Herbelin, "The Role of Electron Spin in the NF System," to be published.

^{**} M. A. Kwok, J. M. Herbelin, and N. Cohen "Collisional Quenching and Radiative Decay Studies of NF($a^{1}\Delta$) and NF($b^{1}\Sigma^{+}$)," to be published.

Munson A. Kwok and Roger L. Wilkins, J. Chem. Phys. 63, 2453 (1975).

²J. M. Herbelin and N. Cohen, Chem. Phys. Lett. <u>20</u>, 605 (1973).

reacts with HI to produce mainly $I(^2P_{3/2})$ atoms.^{3,4} In order to generate sufficient $I(^2P_{1/2})$, it is necessary to pass O_2 through a 2.45 GHz microwave cavity some distance upstream of the reaction zone to yield $O_2(a^1\Delta)$. The O atoms are suppressed by the heated mercuric oxide coating in the quartz sidearm. The transfer $O_2(a^1\Delta) + I(^2P_{3/2})$ is known to be extremely rapid for the efficient production of $I(^2P_{1/2})$.

Flow conditions are chosen such that $O_2(a^1\Delta)$ is excess to I for ease of analysis, but not in sufficient excess to obliterate evidence of the I(2P1/2) + $NF(a^{1}\Delta)$ transfer by the alternative process $O_{2}(a^{1}\Delta) + NF(a^{1}\Delta)$. Typical reagent flowrates for Case 4 in Fig. 1 would be in µmol-sec⁻¹: F, 15; NF₂, 0.15; O₂, 50; D₂, 150; HI, 0.08. Experience^{6,7} indicates that the amount of $O_2(a^1\Delta)$ generated in the usual 2.45 GHz 70 W microwave is 5% of O_2 . The tube operates at 1 Torr, 298 K and 6800 cm sec⁻¹. Case 1 is a previously used baseline. 2 Cases 2 and 3 verify the necessity of the simultaneous presence, in Case 4, of $O_2(a^1\Delta)$ and I for significant pumping of NF($b^1\Sigma^+$). The axial number density plot of NF($b^1\Sigma^+$) versus Z shown in Fig. 1 for Case 4 shows this strong pumping. We interpret Case 4 to mean that I(2P_{1/2}) is rapidly produced with the presence of $O_2(a^2\Delta)$ and then rapid transfer occurs, as described by Eq. (1). Axial number density plots of NF($a^{1}\Delta$) for Cases 1 through 4 are virtually identical. NF($a^{1}\Delta$) behaves as a reservoir because, in all cases, the NF($b^1\Sigma^+$) is at most only 1% of observed NF($a^1\Delta$) concentrations.

A quantitative estimate of the transfer rate coefficient for Eq. (1) can be made. The growth of NF($b^1\Sigma^+$) for Case 4 in Fig. 1 is found to be quadratic in Z (or time). This implies that either NF($a^1\Delta$) or I($^2P_{1/2}$) is in a

³W. E. Jones, S. D. MacKnight, and Lo Teng, Chem. Rev. 73, 407 (1973).

⁴P. Cadman and J. C. Polanyi, J. Phys. Chem. <u>72</u>, 3715 (1968).

⁵R. G. Derwent and B. A. Thrush, Chem. Soc. Faraday Disc. <u>53</u>, 162 (1972).

⁶R. G. Derwent and B. A. Thrush, Trans. Faraday Soc. <u>67</u>, 2036 (1971).

 $^{^{7}}$ L. W. Bader and E. A. Ogryzlo, Disc. Faraday Soc. 37, 46 (1964).




Figure 1. Axial Variations of the Number Density of NF(b¹Σ⁺), Averaged Along the Tube Diameter. Case 1: F, NF₂, D₂, O₂. Case 2: F, NF₂, D₂, O₂, O₂(a¹Δ). Case 3: F, NF₂, D₂, O₂, HI. Case 4: F, NF₂, D₂, O₂, O₂(a¹Δ), HI. The solid lines represent model fits.

linear region of growth. A simple five-kinetic equation analysis for the one unknown rate shows that the lower bound for k_1 , in Eq. (1), is 2.6×10^{14} cm 3 mol $^{-1}$ sec $^{-1}$. The estimate is a bound because the absolute $O_2(a^1\Delta)$ density has not been measured but has been assumed to be $\leq 5\%$ of O_2 , and k_1 is inversely proportional to $O_2(a^1\Delta)$. A more detailed description of this study will be given elsewhere.

LABORATORY OPERATIONS

The Laboratory Operations of The Aerospace Corporation is conducting experimental and theoretical investigations necessary for the evaluation and application of scientific advances to new military concepts and systems. Versatility and flexibility have been developed to a high degree by the laboratory personnel in dealing with the many problems encountered in the nation's rapidly developing space and missile systems. Expertise in the latest scientific developments is vital to the accomplishment of tasks related to these problems. The laboratories that contribute to this research are:

Aerophysics Laboratory: Launch and reentry aerodynamics, heat transfer, reentry physics, chemical kinetics, structural mechanics, flight dynamics, atmospheric pollution, and high-power gas issers.

Chemistry and Physics Laboratory: Atmospheric reactions and atmospheric optics, chemical reactions in polluted atmospheres, chemical reactions of excited species in rocke: plumes, chemical thermodynamics, plasma and laser-induced reactions, laser chemistry, propulsion chemistry, space vacuum and radiation effects on materials, inbrication and surface phenomena, photosinitive materials and sensors, high precision leser ranging, and the application of physics and chemistry to problems of law enforcement and biomedicine.

Electronics Research Laboratory: Electromagnetic theory, devices, and propagation phenomena, including plasma electromagnetics; quantum electronics, lasers, and electro-optics; communication sciences, applied electronics, semi-conducting, superconducting, and crystal device physics, optical and acoustical imaging; atmospheric pollution, millimeter wave and far-infrared technology.

Materials Sciences Laboratory: Development of new materials; metal matrix composites and new forms of carbon; test and evaluation of graphite and ceramics in reentry; spacecraft materials and electronic components in nuclear weapons environment; application of fracture mechanics to stress corrosion and fatigue-induced fractures in structural metals.

Space Sciences Laboratory: Atmospheric and ionospheric physics, radiation from the atmosphere, density and composition of the atmosphere, aurorae and airglow; magnetospheric physics, cosmic rays, generation and propagation of plasma waves in the magnetosphere; solar physics, studies of solar magnetic fields: space astronomy, x-ray astronomy; the effects of nuclear explosions, magnetic storms, and solar activity on the earth's atmosphere, ionosphere, and magnetosphere; the effects of optical, electromagnetic, and particulate radiations in space on space systems.

THE AEROSPACE CORPORATION El Segundo, California